

COSMOLOGY OF ROLLING TACHYON *

CHANJU KIM

Department of Physics, Ewha Womans University, Seoul 120-750, Korea
E-mail: cjkim@ewha.ac.kr

HANG BAE KIM

*Institute of Theoretical Physics, University of Lausanne,
CH-1015 Lausanne, Switzerland*
E-mail: HangBae.Kim@ipt.unil.ch

YOONBAI KIM AND O-KAB KWON

*BK21 Physics Research Division and Institute of Basic Science,
Sungkyunkwan University, Suwon 440-746, Korea*
and
*School of Physics, Korea Institute for Advanced Study,
207-43, Cheongryangri-Dong, Dongdaemun-Gu, Seoul 130-012, Korea*
E-mail: yoonbai@skku.ac.kr, okwon@newton.skku.ac.kr

We study dynamics of rolling tachyon and Abelian gauge field on unstable D-branes, of which effective action is given by Born-Infeld type nonlocal action. Possible cosmological evolutions are also discussed. In the Einstein frame of string cosmology, every expanding flat universe is proven to be decelerating.

1 Introduction

Recent analysis of string theory ^{1,2,3} suggests that classical dynamics of tachyon on an unstable D-brane is described by a scalar Born-Infeld type action with a runaway potential ⁴. Characteristic of such tachyon effective theory is absence of perturbative tachyonic modes around the potential minimum and existence of spatially homogeneous rolling tachyon configuration ⁵. In the background of the rolling tachyon, Abelian gauge field on the D-brane cannot support perturbative spectrum at late-time ^{6,7,9}. In this review, we summarize firstly bosonic effective action both in the bulk and on the D-brane, together with shapes of the tachyon potential. In section 3, we introduce Carrollian limit, and then discuss nonexistence of perturbative spectra of the tachyon and the Abelian gauge field in the potential minimum, of which dynamics are governed by nonlocal D-brane action. For the tachyon, there exists the classical rolling tachyon solution connecting smoothly unstable maximum

*PROCEEDINGS OF PAC MEMORIAL SYMPOSIUM ON THEORETICAL PHYSICS, PP.209-239 (CHUNGBUM PUBLISHING HOUSE, SEOUL). TALK WAS GIVEN BY Y. KIM.

of the potential and stable minimum at infinity.

Though the situation seems not to be matured yet, string cosmology must be an intriguing subject to be tackled at every step of progress in both string theory and observational cosmology¹⁰. Recently, inspired by string theory D-branes and heterotic M theory, brane cosmology attracted much attention. Brane cosmology assumes that our universe starts out with branes embedded in the higher dimensional spacetime, either stable or unstable. In such context, cosmology of the rolling tachyon must be an attractive subject^{11,12,13,14,15}. Topics included inflation, dark matter, cosmological perturbation, and reheating despite stringent difficulty in the simplest versions of this theory. A nice aspect for the cosmology is nothing but the absence of open string excitations so that classical analysis of late-time based on homogeneity and isotropy may lead to a solid prediction. In section 4, we briefly describe system of gravity and the tachyon. The obtained universe in the simplest set predicts that expanding universe is accelerating at onset stage but turns to decelerating phase at late-time¹¹. Finally, the scale factor approaches a constant. Our main concern is string cosmology involving graviton, dilaton, and the tachyon. In the string frame, we obtain flat space solutions at both the early epochs and late-times in closed form. When transformed to the Einstein frame, analysis shows that every expanding universe of the graviton-tachyon-dilaton system should be decelerating irrespective of the specific shape of the tachyon potential. Specific form of the scale factor is proportional to (time)^{1/2}.

We conclude with a summary of this paper. Sometimes, same notations are used in different sections for convenience.

2 Setup

Effective action of bosonic sector of D3-brane system of tension T_3 is described by the following action in string frame^{16,1}

$$S = \frac{1}{2\kappa^2} \int d^4x \sqrt{-g} e^{-2\Phi} \left(R + 4\partial_\mu \Phi \partial^\mu \Phi - \frac{1}{12} H_{\mu\nu\rho} H^{\mu\nu\rho} - 2\Lambda \right) - T_3 \int d^4x e^{-\Phi} V(T) \sqrt{-\det(g_{\mu\nu} + \partial_\mu T \partial_\nu T + B_{\mu\nu} + F_{\mu\nu})}, \quad (1)$$

where cosmological constant Λ is assumed to be vanished in what follows.^a The (quadratic) action in the first line of Eq. (1) is bulk action of graviton $g_{\mu\nu}$, dilaton Φ , and antisymmetric tensor field $B_{\mu\nu}$ of rank 2 of which field strength $H_{\mu\nu\rho}$ is defined by $H_{\mu\nu\rho} = \partial_\mu B_{\nu\rho} + \partial_\nu B_{\rho\mu} + \partial_\rho B_{\mu\nu}$. The action induced on the D3-brane is Born-Infeld type nonlocal action given in the second line of

^aOther effective actions different from Eq. (1) can also be considered. For example boundary string field theory(BSFT) action for a non-BPS D-brane was also proposed and studied².

Eq. (1), which involves kinetic terms of tachyon T and Abelian gauge field A_μ satisfying $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$. Note that the tachyon T and the Abelian gauge field A_μ were rescaled as $T \rightarrow \sqrt{2\pi\alpha'}T$ and $A_\mu \rightarrow 2\pi\alpha'A_\mu$ to absorb $2\pi\alpha'$ in front of the kinetic terms.

Since we will deal with two classical topics in this paper, i.e., one is string cosmology including the rolling tachyon¹⁵ and the other is evolution of electromagnetic waves in the background of the rolling tachyon⁹, let us firstly read equations of motion from the action (1)

$$\begin{aligned} R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R = & -\Lambda g_{\mu\nu} - 2[\nabla_\mu \nabla_\nu \Phi + g_{\mu\nu}(\nabla_\alpha \Phi \nabla^\alpha \Phi - \nabla_\alpha \nabla^\alpha \Phi)] \\ & + \left(\frac{1}{4}H_{\mu\alpha\beta}H_\nu^{\alpha\beta} - \frac{1}{24}g_{\mu\nu}H_{\alpha\beta\gamma}H^{\alpha\beta\gamma} \right) \\ & + \kappa^2 T_3 e^\Phi V(T) \left(\frac{g_{\mu\alpha}C^{\alpha\beta}\partial_\beta T \partial_\nu T}{\sqrt{gX}} - g_{\mu\nu}\sqrt{g^{-1}X} \right), \quad (2) \end{aligned}$$

$$4(-\nabla_\mu \nabla^\mu \Phi + \nabla_\mu \Phi \nabla^\mu \Phi) = R - 2\Lambda - \frac{1}{12}H_{\alpha\beta\gamma}H^{\alpha\beta\gamma} - \kappa^2 T_3 e^\Phi V(T)\sqrt{g^{-1}X}, \quad (3)$$

$$\nabla_\mu H^{\mu\nu\lambda} = 2\nabla_\mu \Phi H^{\mu\nu\lambda}, \quad (4)$$

$$\begin{aligned} C_{(s)}^{\mu\nu} \nabla_\mu \nabla_\nu T - \frac{C_{(s)}^{\mu\nu} C^{\alpha\beta} \nabla_\mu T \nabla_\nu X_{\alpha\beta} - X \nabla^\mu T \nabla_\nu C_{(s)}^{\mu\nu}}{2X} - C_{(s)}^{\mu\nu} \nabla_\mu T \nabla_\nu \Phi \\ - \frac{1}{V} \frac{dV}{dT} \left(X - C_{(s)}^{\mu\nu} \nabla_\mu T \nabla_\nu T \right) = 0, \quad (5) \end{aligned}$$

where $X_{\mu\nu} = g_{\mu\nu} + B_{\mu\nu} + F_{\mu\nu} + \partial_\mu T \partial_\nu T$, $X = \det(X_{\mu\nu})$, $C^{\mu\nu}$ is the cofactor of $X_{\mu\nu}$, and $C_{(s)}^{\mu\nu}$ is the symmetric part of $C^{\mu\nu}$.

The tachyon potential $V(T)$ multiplied in front of the Born-Infeld type action (1) is known to measure varying tension of the unstable D-branes. Though there seem no consensus and no exact computation on the form of tachyon potential $V(T)$, it should have its maximum, $\max(V(T)) = 1$, and minimum, $\min(V(T)) = 0$. Some known computations from string theories are as follows⁴. For bosonic string theory, we have

$$V(T) = \begin{cases} \frac{T^2}{8} \exp\left(1 - \frac{T^2}{8}\right) & \text{for small } T \text{ but } T \geq 1 \\ \exp\left(-\frac{T}{2}\right) & \text{for large } T \end{cases}, \quad (6)$$

and, for superstring theory, we have

$$V(T) = \begin{cases} \exp\left(-\frac{T^2}{8\ln 2}\right) & \text{for small } T \text{ but } T \geq 0 \\ \exp\left(-\frac{T}{\sqrt{2}}\right) & \text{for large } T \end{cases}. \quad (7)$$




Figure 1. Tachyon potential $V(T) = \frac{1}{\cosh T}$

String theory calculation (6)–(7) tells us that favorable form of the tachyon potential supports $V \sim e^{-T^2}$ for small T and $V \sim e^{-T}$ for large T . Since the potential $V(T)$ must be a smooth function in the intermediate region, connecting two asymptotic expressions, a typical example is ¹³

$$V(T) = \left(1 + \frac{T}{T_0}\right) \exp\left(-\frac{T}{T_0}\right). \quad (8)$$

In this paper, considering convenience in analytic computation, we adopt

$$V(T) = \frac{1}{\cosh\left(\frac{T}{T_0}\right)} \quad (9)$$

which is also consistent with the results of string theory (6)–(7) (See Fig. 1).

3 Tachyon and Gauge Field

Among various bosonic degrees of freedom, dynamics of the tachyon T and the gauge field A_μ is governed by nonlocal kinetic terms in the brane action (1). Before discussing their dynamics directly, let us consider a massless noninteracting real scalar field ϕ described by Lorentz-invariant linear wave

equation

$$\left(\frac{1}{v^2} \frac{\partial^2}{\partial t^2} - \nabla^2\right) \phi = 0. \quad (10)$$

If we take two singular limits known as the Inönö-Wigner limits, infinite velocity limit, $v \rightarrow \infty$, goes to Galileo's limit of action at a distance described by an elliptic equation :

$$\nabla^2 \phi = 0 \quad (11)$$

which we encounter frequently in Newtonian dynamics. On the other hand, the opposite velocity dominated limit, $v \rightarrow 0$, called as Carrollian limit is described by an ordinary differential equation :

$$\frac{d^2}{dt^2} \phi = 0. \quad (12)$$

A natural implication from the above discussion is possible absence of plane waves in the Carrollian limit of corresponding field theory. Instead time evolution of homogeneous configuration may be dominated. Though both limits are singular in the context of group theory, we will use the speed v as a continuous parameter of range $(0 \leq v \leq 1)$ as we have done in the Galileo's limit where its value is taken from one to infinity.

In this section, we will briefly review rolling tachyon in flat spacetime with disappearance of perturbative tachyonic spectrum at the bottom of the potential $V(T)$ and discuss singular evolution of perturbative electromagnetic field in background of the rolling tachyon.

3.1 Rolling tachyon in flat spacetime

Let us begin this subsection with discussing physics of unstable D-branes in the simplest setup. If we consider solely motion of the tachyon in flat spacetime, it is depicted by ⁵

$$S_T = -T_3 \int d^4x V(T) \sqrt{1 + \partial_\mu T \partial^\mu T}, \quad (13)$$

where specific form of the tachyon potential $V(T)$ is chosen by Eq. (9). Note that $g^{-1} \det(g_{\mu\nu} + \partial_\mu T \partial_\nu T) = 1 + \partial_\mu T \partial^\mu T$ up to the quadratic terms and it holds in an exact form as far as homogeneous configurations are concerned such as the cosmology described in terms of Robertson-Walker metric of our interest. Equation of motion is

$$\frac{1}{\sqrt{-\eta}} \partial_\mu (V \sqrt{-\eta} \eta^{\mu\nu} \partial_\nu T) = \frac{dV}{dT}, \quad (14)$$

where $\bar{\eta} = -(1 + \partial_\mu T \partial^\mu T)$ and

$$\bar{\eta}_{\mu\nu} = \eta_{\mu\nu} + \partial_\mu T \partial_\nu T, \quad \bar{\eta}^{\mu\nu} = \eta^{\mu\nu} - \frac{\partial^\mu T \partial^\nu T}{1 + \partial_\rho T \partial^\rho T}. \quad (15)$$

Energy-momentum tensor $T_{\mu\nu}$ is given by

$$T_{\mu\nu} = -T_3 V(T) \eta_{\mu\rho} \eta_{\nu\sigma} \sqrt{-\bar{\eta}} \bar{\eta}^{\rho\sigma}. \quad (16)$$

Since $T = 0$ is unique stationary point of the tachyon potential $V(T)$, we perturb the action (13) up to quadratic terms to read perturbative spectrum. For small T , we have

$$S_T + T_3 \int d^4x \approx T_3 \int d^4x \left(-\frac{1}{2} \partial_\mu T \partial^\mu T + \frac{1}{2T_0^2} T^2 \right), \quad (17)$$

where the positive cosmological constant at $T = 0$ is subtracted in the action. As expected, the sign in front of mass term is negative so it is tachyonic on top of the potential $V(T = 0)$. For large T , the tachyon potential is approximated as $V(T) \approx 2e^{-T/T_0}$. Introducing a new small order parameter such as $\phi = 2\sqrt{2}T_0 e^{-T/2T_0}$, we obtain a quadratic action with positive mass term

$$S_T \approx T_3 \int d^4x \left[-\frac{1}{2} \partial_\mu \phi \partial^\mu \phi - \frac{1}{2} \left(\frac{1}{\sqrt{2}T_0} \right)^2 \phi^2 \right]. \quad (18)$$

Though this seems to imply existence of a perturbative tachyonic spectrum of mass $1/\sqrt{2}T_0$, it is an artifact of quadratic approximation since $T \rightarrow \infty$ is not a stationary point of the tachyon potential as shown in the Fig. 1. To be specific, let us consider a plane wave solution $\phi \sim e^{ik_\mu x^\mu}$ and then we have $\partial_\mu T \sim -2iT_0 k_\mu$. Inserting these into the equation of motion (14), we get the following dispersion relation

$$\frac{1}{\sqrt{1 - 4T_0^2 k^2}} \frac{d \ln V}{dT} = 0. \quad (19)$$

Note that $d(\ln V)/dT|_{T \rightarrow \infty} = -T_0 \neq 0$, the dispersion relation does not have any nontrivial solution but can only be satisfied for infinite k^2 . The mass-shell condition we derived through a quadratic approximation (18) was reproduced by a Taylor expansion of Eq. (19) up to quadratic terms for small k^2 .

Absence of perturbative tachyon spectrum does not lead to nonexistence of classical configurations. We may take into account an intriguing lesson from the Carrollian limit (12) of the wave equation. Suppose that we have a homogeneous tachyon configuration with $\partial_i T = 0$ and consider its classical time evolution $T(t)$. Then only diagonal components of the energy-momentum tensor (16) survive and the equation of motion (14) is integrated to conservation

of energy $\rho_T \equiv T_{00}$, i.e., Hamiltonian density $\mathcal{H}_T$ is

$$\mathcal{H}_T = \rho_T = T_3 \frac{V(T)}{\sqrt{1 - \dot{T}^2}} = T_3 \epsilon, \quad (20)$$

where $\epsilon = V(T_i)/\sqrt{1 - \dot{T}_i^2} \geq 1$ and $T_i, \dot{T}_i$ are the initial values of T and $\dot{T}$. It is analogous to the aforementioned Carrollian limit (12). Constancy of the Hamiltonian density (20) forces $\dot{T} \rightarrow 1$ for any finite ϵ as $T \rightarrow \infty$. For the tachyon potential (9), we have a solution of rolling tachyon

$$T(t) = T_0 \sinh^{-1} \left(a e^{t/T_0} - \frac{b}{4a} e^{-t/T_0} \right), \quad (21)$$

where $a = \frac{1}{2} \left[\sinh(T_i/T_0) + \sqrt{\sinh^2(T_i/T_0) + b} \right]$ and $b = (\epsilon^2 - 1)/\epsilon^2$. At late-time regime, the obtained tachyon solution (21) is approximated as expected, i.e., $T(t) \approx t$ irrespective of the initial values of T and $\dot{T}$. Note that the pressure $p_T \equiv T_i/3$ is negative and approaches zero as time elapses, which coincides with vanishing Lagrangian limit

$$p_T = \mathcal{L}_T = -T_3 V(T) \sqrt{1 - \dot{T}^2} \xrightarrow{t \rightarrow \infty} 0. \quad (22)$$

From equation of state $p_T = w_T \rho_T$, we read

$$w_T = -(1 - \dot{T}^2) = -\frac{V(T(t))^2}{\epsilon^2} \leq 0, \quad (23)$$

where $w_T = -1/\epsilon^2$ at $T = 0$ and $w_T \rightarrow 0$ at $T \rightarrow \infty$.

3.2 Gauge field and rolling tachyon

When we turn off antisymmetric tensor field of second rank $B_{\mu\nu}$ in the Born-Infeld type action in Eq. (1), effective bosonic action of unstable D3-brane system is given by ^{6,7,3,8,9}

$$S = -T_3 \int d^4x V(T) \sqrt{-\det(\eta_{\mu\nu} + \partial_\mu T \partial_\nu T + F_{\mu\nu})}. \quad (24)$$

To proceed, we introduce a few notations. We first define

$$X_{\mu\nu} \equiv \eta_{\mu\nu} + \partial_\mu T \partial_\nu T + F_{\mu\nu}, \quad (25)$$

$$X \equiv \det(X_{\mu\nu}), \quad (26)$$

and barred covariant field strength tensor $\bar{F}_{\mu\nu}$

$$\bar{F}_{\mu\nu} = F_{\mu\nu} \quad (27)$$

with Eq. (15). Then we have contravariant barred field strength tensor $\bar{F}^{\mu\nu}$ and its dual field strength $\bar{F}_{\mu\nu}^*$

$$\bar{F}^{\mu\nu} = \bar{\eta}^{\mu\alpha} \bar{\eta}^{\nu\beta} F_{\alpha\beta}, \quad \bar{F}_{\mu\nu}^* = \frac{\bar{\epsilon}_{\mu\nu\alpha\beta}}{2} \bar{F}^{\alpha\beta} = \frac{\bar{\epsilon}_{\mu\nu\alpha\beta}}{2} \bar{\eta}^{\alpha\gamma} \bar{\eta}^{\beta\delta} F_{\gamma\delta}, \quad (28)$$

where $\bar{\epsilon}_{\mu\nu\alpha\beta} = \sqrt{-\bar{\eta}} \epsilon_{\mu\nu\alpha\beta}$ with $\epsilon_{0123} = 1$.

In terms of barred quantities Eq. (26) is computed as

$$X = \bar{\eta} \left[1 + \frac{1}{2} \bar{F}_{\mu\nu} \bar{F}^{\mu\nu} - \frac{1}{16} (\bar{F}_{\mu\nu}^* \bar{F}^{\mu\nu})^2 \right]. \quad (29)$$

Then equations of motion for the tachyon T and the gauge field A_μ are

$$\partial_\mu \left(\frac{V}{\sqrt{-X}} C_S^{\mu\nu} \partial_\nu T \right) + \sqrt{-X} \frac{dV}{dT} = 0, \quad (30)$$

$$\partial_\mu \left(\frac{V}{\sqrt{-X}} C_A^{\mu\nu} \right) = 0. \quad (31)$$

Here $C_S^{\mu\nu}$ and $C_A^{\mu\nu}$ are symmetric and asymmetric part, respectively, of the cofactor,

$$C^{\mu\nu} = \bar{\eta} (\bar{\eta}^{\mu\nu} + \bar{F}^{\mu\nu} + \bar{\eta}^{\mu\alpha} \bar{\eta}^{\beta\gamma} \bar{\eta}^{\delta\nu} \bar{F}_{\alpha\beta}^* \bar{F}_{\gamma\delta} + \bar{\eta}^{\mu\alpha} \bar{\eta}^{\beta\gamma} \bar{F}_{\alpha\beta}^* \bar{F}_{\gamma\delta}^* \bar{F}^{\delta\nu}), \quad (32)$$

namely,

$$C_S^{\mu\nu} = \bar{\eta} (\bar{\eta}^{\mu\nu} + \bar{\eta}^{\mu\alpha} \bar{\eta}^{\beta\gamma} \bar{\eta}^{\delta\nu} \bar{F}_{\alpha\beta}^* \bar{F}_{\gamma\delta}), \quad (33)$$

$$C_A^{\mu\nu} = \bar{\eta} (\bar{F}^{\mu\nu} + \bar{\eta}^{\mu\alpha} \bar{\eta}^{\beta\gamma} \bar{F}_{\alpha\beta}^* \bar{F}_{\gamma\delta}^* \bar{F}^{\delta\nu}). \quad (34)$$

Energy-momentum tensor $T_{\mu\nu}$ is given by

$$T_{\mu\nu} = -\frac{T_3 V(T)}{\sqrt{-X}} [-\eta_{\mu\nu} X + C_{\mu\rho} (\partial_\nu T \partial^\rho T + F_\nu{}^\rho)], \quad (35)$$

where $C_{\mu\nu} \equiv \eta_{\mu\alpha} \eta_{\nu\beta} C^{\alpha\beta}$. The diagonal components can be identified as density and pressure,

$$\rho = -\frac{T_3 V(T)}{\sqrt{-X}} [X + C_{0\mu} (\dot{T} \partial^\mu T + F_0{}^\mu)], \quad (36)$$

$$p = -\frac{T_3 V(T)}{3\sqrt{-X}} [-3X + C_{0\mu} (\dot{T} \partial^\mu T + F_0{}^\mu) + C^{\mu\nu} (\partial_\mu T \partial_\nu T + F_{\mu\nu})]. \quad (37)$$

From equation of state $p = w\rho$, we read

$$w = \frac{1}{3} - \frac{4X - C^{\mu\nu} (\partial_\mu T \partial_\nu T + F_{\mu\nu})}{3 [X + C_{0\mu} (\dot{T} \partial^\mu T + F_0{}^\mu)]}. \quad (38)$$

Thus the trace of the total energy-momentum tensor is given by

$$T^\mu{}_\mu = -\frac{T_3 V(T)}{\sqrt{-X}} \left[(4 + 3\partial_\rho T \partial^\rho T) + \left((1 + \frac{1}{2}\partial_\rho T \partial^\rho T) \delta^\mu{}_\sigma - \partial^\mu T \partial_\sigma T \right) F_{\mu\nu} F^{\sigma\nu} \right]. \quad (39)$$

The equations of motion (30) and (31) admit homogeneous solutions $T(t)$ and $A_\mu(t)$ which satisfy

$$\begin{aligned} \partial_0 \left(\frac{V}{\sqrt{-X}} \dot{T} \right) + \sqrt{-X} \frac{dV}{dT} &= 0, \\ \partial_0 \left(\frac{V}{\sqrt{-X}} E^i \right) &= 0, \end{aligned} \quad (40)$$

where $X = -1 + \dot{T}^2 + \mathbf{E}^2$. Solving these equations we find that $V/\sqrt{-X} = \text{constant}$ and $\mathbf{E} = \text{constant}$. The first equation is nothing but the statement of conservation of energy.

When $\mathbf{E} = 0$, the solution reduces to that in the theory with pure tachyon field given in the subsection 3.1. When $\mathbf{E} \neq 0$, one can simply read the solution from Eq. (21) by rescaling $\epsilon^2 \rightarrow \epsilon^2(1 - E^2)$ and $T_0 \rightarrow T_0/\sqrt{1 - E^2}$ in the exponents of Eq. (21).

Now we consider the propagation of the gauge field A_μ in the rolling tachyon background (21). For this purpose we linearize the equation of motion (31) in this background. Written in terms of electric and magnetic fields, Ampère's law is modified as

$$\nabla \times \mathbf{B} = \frac{1}{V\sqrt{1 - \dot{T}^2}} \frac{\partial}{\partial t} \left(\frac{V}{\sqrt{1 - \dot{T}^2}} \mathbf{E} \right). \quad (41)$$

Combined with the Bianchi identity, the wave equations of the electric field $\mathbf{E}$ and the magnetic field $\mathbf{B}$ become

$$\frac{1}{V\sqrt{1 - \dot{T}^2}} \frac{\partial}{\partial t} \left(\frac{V}{\sqrt{1 - \dot{T}^2}} \frac{\partial \mathbf{B}}{\partial t} \right) - \nabla^2 \mathbf{B} = 0, \quad (42)$$

$$\frac{\partial}{\partial t} \left[\frac{1}{V\sqrt{1 - \dot{T}^2}} \frac{\partial}{\partial t} \left(\frac{V\mathbf{E}}{\sqrt{1 - \dot{T}^2}} \right) \right] - \nabla^2 \mathbf{E} = 0. \quad (43)$$

Note that these equations are not identical. Since the Hamiltonian density $\mathcal{H}_T$ (20) is a constant of motion, the wave equations (42) and (43) reduce to

$$\frac{\epsilon^2}{V^2} \frac{\partial^2 \mathbf{B}}{\partial t^2} - \nabla^2 \mathbf{B} = 0, \quad (44)$$

$$\frac{\epsilon^2}{V^2} \frac{\partial^2 \mathbf{E}}{\partial t^2} + \frac{d}{dt} \left(\frac{\epsilon^2}{V^2} \right) \frac{\partial \mathbf{E}}{\partial t} - \nabla^2 \mathbf{E} = 0. \quad (45)$$

One can also compute the energy density (36) and pressure (37) as are given by

$$\rho_{\text{em}} \equiv T_{00} - T_{00}|_{F_{\mu\nu}=0} \approx \frac{T_3}{2} \frac{V(T)}{\sqrt{1-\dot{T}^2}} \left(\frac{\mathbf{E}^2}{1-\dot{T}^2} + \mathbf{B}^2 \right), \quad (46)$$

$$p_{\text{em}} \equiv \frac{1}{3}(T_{ii} - T_{ii}|_{F_{\mu\nu}=0}) \approx \frac{T_3}{6} \frac{V(T)}{\sqrt{1-\dot{T}^2}} \left[\mathbf{E}^2 + (1-\dot{T}^2)\mathbf{B}^2 \right]. \quad (47)$$

Similarly, we read w from Eq. (38)

$$w_{\text{em}} \equiv \frac{p_{\text{em}}}{\rho_{\text{em}}} \approx \frac{1}{3}(1-\dot{T}^2). \quad (48)$$

The trace of the energy-momentum tensor (39) becomes

$$T_{em}{}^\mu{}_\mu \equiv T^\mu{}_\mu - T^\mu{}_\mu|_{F_{\mu\nu}=0} \approx -\frac{T_3}{2} \frac{V(T)}{\sqrt{1-\dot{T}^2}} \dot{T}^2 \left(\frac{\mathbf{E}^2}{1-\dot{T}^2} + \mathbf{B}^2 \right). \quad (49)$$

The speed of wave propagation is altered by $V(T(t))$ and thus time-varying. One might expect from Eq. (44) that the propagation speed roughly follows $V(t)$, but this is not necessarily so as we will see below. Considering the shape of the potential in Eqs. (6) and (7), we see that it is the light speed on the top of the potential and decreases monotonically to zero as the rolling of the tachyon proceeds. For illustrative purpose, we take the tachyon potential (8) and the solution (21). Inserting the solution into the tachyon potential, we have an expression of V as a function of time t

$$V(T(t)) = \frac{1}{\sqrt{1 + \left(ae^{t/T_0} - \frac{b}{4a} e^{-t/T_0} \right)^2}}. \quad (50)$$

As shown in Fig. 2, $V(T(t))$ changes in the scale of a few T_0 's from the light speed to zero, which is nothing but elapsed time from the onset of the tachyon rolling to the late-time. Since the electric field and the magnetic field obey different wave equations, they show different behaviors in propagation.

To investigate in detail the propagation of electromagnetic wave in the rolling tachyon background, let us try to find a plane-wave solution of the wave equations with a time-varying propagation speed, (44) and (45). We consider the plane-wave propagating along the x -axis, $\mathbf{k} = k\hat{\mathbf{x}}$, and make an ansatz $\mathbf{B} = B_z(t, x)\hat{\mathbf{z}}$ with

$$B_z(t, x) = B(t)e^{ik[x-W_B(t)]}, \quad (51)$$

where $B(t)$ and $W_B(t)$ are real. We assume that $V(0) \approx 1$ and $\dot{W}_B(0) \approx 1$ so that the ansatz (51) gives the normal plane-wave initially. Since $\mathbf{E}$ and $\mathbf{B}$ are related through the Ampère's law (41) and Faraday's law

$$\nabla \times \mathbf{E} + \frac{\partial \mathbf{B}}{\partial t} = 0, \quad (52)$$




Figure 2. (Top) tachyon potential as a function of time, Eq. (50) for $a = 0.1$ and $b = 0$ ($\epsilon = 1$). (Middle) amplitudes and (bottom) propagation speeds of magnetic field (upper three lines) and electric field (lower three lines) for $kT_0 = 0.1$ (dashed line), $kT_0 = 1$ (solid line), and $kT_0 = 10$ (dotted line).

the electric field $\mathbf{E} = E_y(t, x)\hat{\mathbf{y}}$ is then given by

$$E_y(t, x) = E(t)e^{ik[x - W_E(t)]}, \quad (53)$$

with

$$E = \sqrt{\dot{W}_B^2 B^2 + k^{-2} \dot{B}^2}, \quad (54)$$

$$W_E = W_B - k^{-1} \tan^{-1} \left(\frac{k^{-1} \dot{B}}{\dot{W}_B B} \right). \quad (55)$$

With this ansatz, Eq. (44) reduces to

$$\ddot{B} - k^2 \left(\dot{W}_B^2 - \epsilon^{-2} V^2 \right) B = 0, \quad (56)$$

$$2\dot{B}\dot{W}_B + B\ddot{W}_B = 0. \quad (57)$$

Eq. (57) implies

$$\dot{W}_B = \frac{1}{\epsilon} \left(\frac{B_0}{B} \right)^2, \quad (58)$$

where B_0 is a constant. From Eq. (55) we also find

$$\dot{W}_E = \frac{V^2}{\epsilon^3} \left(\frac{B_0}{E} \right)^2. \quad (59)$$

Inserting Eq. (58) into Eq. (56), we obtain a second-order nonlinear equation for amplitude of the magnetic field

$$\ddot{B} - \frac{k^2}{\epsilon^2} \left[\left(\frac{B_0}{B} \right)^4 - V^2 \right] B = 0. \quad (60)$$

With the knowledge of $V(T(t))$, like the one given in Eq. (50), we can solve these equations for $B(t)$ and $E(t)$. For $V(T(t))$ of Eq. (50), the numerical solution is shown in Fig. 2. Let us first examine a few limiting cases.

On the top of the potential $V = 1$, we get a normal plane-wave solution $\dot{W}_B = \dot{W}_E = 1/\epsilon$ with amplitudes of $B = B_0$ and $E = B_0/\epsilon$ as expected from the original wave equations (44)–(45). At the top of the potential, the equation of states (48) gives $w_{\text{em}} = 1/3$ and the trace of the energy-momentum tensor $T_{\text{em}}^\mu{}_\mu$ up to quadratic terms (49) vanishes.

Once the tachyon starts rolling from the top of the potential, the fluctuations of the gauge fields starts deviating from the normal plane wave solution. With the initial condition $V(t=0) = 1$ for which $4a^2 = b$, $V(T(t))$ in Eq. (50) reduces to

$$V(T(t)) = \frac{1}{\sqrt{1 + b \sinh^2 t/T_0}}. \quad (61)$$

Assuming that $b = (\epsilon^2 - 1)/\epsilon^2$ is small, which means the tachyon rolls slowly initially, we can solve Eqs. (60) and (58) to the first order in b :

$$\begin{aligned} B(t) &= B_0 \left[1 + \frac{bk^2T_0^2}{8(1+k^2T_0^2)} \sinh^2 t/T_0 \right], \\ W_B(t) &= \frac{t}{\epsilon} + \frac{b}{4\epsilon} \left[t - \frac{k^2T_0^3}{2(1+k^2T_0^2)} \sinh 2t/T_0 \right], \end{aligned} \quad (62)$$

where we rescaled B_0 so that $B(0) = B_0$. This solution is valid as long as $be^{t/T_0} \ll 1$. Thus the amplitude initially grows quadratically as the tachyon starts rolling. On the other hand, electric field can be calculated from (54) and (59),

$$\begin{aligned} E(t) &= B_0 \left[1 - \frac{b}{4(1+k^2T_0^2)} (1 + 2k^2T_0^2 + k^2T_0^2 \sinh^2 t/T_0) \right], \\ W_E(t) &= \frac{t}{\epsilon} - \frac{b}{8\epsilon(1+k^2T_0^2)} [2k^2T_0^2t + (2 + k^2T_0^2) \sinh 2t/T_0]. \end{aligned} \quad (63)$$

Note that $E(t)$ and $\dot{W}_E$ tends to decrease in contrast with the behavior of magnetic field. This can be confirmed in Fig. 2. As time elapses, w_{em} in Eq. (48) starts decreasing but $T_{\text{em}}{}^\mu{}_\mu$ up to quadratic terms in Eq. (49) develops nonvanishing contributions.

At late-time, the value of the potential (50) decays exponentially to zero

$$V(T(t)) \approx \frac{1}{\sqrt{b}e^{t/T_0}}. \quad (64)$$

In this case, it is more convenient to work with the original equation (44) from which we see that the time dependence is exponentially suppressed at late time. Therefore the magnetic field linearly diverges in general,

$$B(t) = B_1t + B_2 + O(e^{-t/T_0}). \quad (65)$$

Then the propagation speed decreases according to Eq. (58) (See also Fig. 2). Since B linearly increases, our quadratic approximation of the Born-Infeld action is not valid as t gets large. However, one may still extract some relevant physics as the tachyon rolls to the vacuum, $V \rightarrow 0$. The electromagnetic waves become frozen and stop propagating. For the magnetic field, short wavelengths freeze earlier than long wavelengths. For the electric field in this limit, we have from Eq. (52),

$$E(t) = B_1/k, \quad (66)$$

which is a constant. Note that in (59) $\dot{W}_E$ contains extra V^2 compared with $\dot{W}_B$ and hence the electric field is frozen more quickly than the magnetic field.

The density ρ_{em} (46) diverges exponentially from the electric component but the pressure p_{em} (47) approaches a finite value, so that w_{em} in Eq. (48)

becomes zero. Similarly, the trace of the energy-momentum tensor (49) also diverges exponentially. Since the Hamiltonian density $\mathcal{H}_T$ for the pure tachyon is almost a constant, the energy density (46) can be compared with that in the materials

$$\frac{T_3}{2} (\mathbf{D} \cdot \mathbf{E} + \mathbf{H} \cdot \mathbf{B}). \quad (67)$$

Then this material may be approximated by a dielectric material at least formally

$$\mathbf{D} \approx \frac{\mathbf{E}}{V(T)^2}, \quad \mathbf{H} \approx \mathbf{B}, \quad (68)$$

where its time-dependent dielectric function diverges as time elapses, $\varepsilon(t) \approx 1/V(T)^2 \xrightarrow{t \rightarrow \infty} \infty$.

4 Cosmological Aspects of Rolling Tachyon

An attractable topic of the rolling tachyon is its application to cosmology^{11,12,13,14,15} as indicated from the beginning⁵. The simplest set for the cosmology of the rolling tachyon is composed of the graviton $g_{\mu\nu}$ and the tachyon T . Therefore, the action is read from Eq. (1)

$$S = \int d^4x \sqrt{-g} \left(\frac{1}{2\kappa^2} R - T_3 V(T) \sqrt{1 + g^{\mu\nu} \partial_\mu T \partial_\nu T} \right). \quad (69)$$

Without any computation, we read a few intriguing subjects from characters of the tachyon potential in Eqs. (6)–(9) (Refer also to Fig. 1). At initial stage of the rolling tachyon, there exists a cosmological constant $V(T \approx 0) \approx 1$ so that one expects easily solutions of inflationary universe. At late-time of $T \rightarrow \infty$, we have tiny nonvanishing but monotonically-decreasing cosmological constant at each instant, which lets us consider it as a possible source of quintessence. Bad news for comprising a realistic cosmological model may be absence of reheating and difficulty of density perturbation at late-time due to disappearance of perturbative degrees reflecting nonexistence of stationary vacuum point in the monotonically-decreasing tachyon potential.

For cosmological solutions, we try a spatially-homogeneous but time-dependent solution

$$ds^2 = -dt^2 + a^2(t) d\Omega_k^2, \quad T = T(t), \quad (70)$$

where $d\Omega_k^2$ corresponds, at least locally, to the metric of S^3 , E^3 , or H^3 according to the value of $k = 1, 0, -1$, respectively. Then the Einstein equations are summarized as

$$\frac{\dot{a}^2}{a^2} + \frac{k}{a^2} = \frac{\kappa^2}{3} \frac{T_3 V(T)}{\sqrt{1 - \dot{T}^2}}, \quad (71)$$

$$\frac{\ddot{a}}{a} = \frac{\kappa^2}{3} \frac{T_3 V(T)}{\sqrt{1 - \dot{T}^2}} \left(1 - \frac{3}{2} \dot{T}^2 \right), \quad (72)$$

and the tachyon equation becomes

$$\frac{\ddot{T}}{1 - \dot{T}^2} + 3 \frac{\dot{a}}{a} \dot{T} + \frac{1}{V} \frac{dV}{dT} = 0, \quad (73)$$

where is equivalent to conservation equation of the energy-momentum tensor.

For the flat universe of $k = 0$, cosmological evolution is rather simple since both sides of Eq. (71) are positive semi-definite. The second equation (72) tells us that the expanding universe is accelerating at onset of $T(t) \approx 0$ since the right-hand side of Eq. (72) is also positive definite. As time t goes, $\dot{T}$ exceeds $\sqrt{2/3}$ so that expansion of the universe slows down and then finally the scale factor $a(t)$ will halt as $a(t) \xrightarrow{t \rightarrow \infty} \text{constant}$. This makes the second term of the tachyon equation (73) vanish at infinite time so that we confirm $\ddot{T} \rightarrow 0$ and $\dot{T} \rightarrow 1$ as $T \rightarrow \infty$.

5 String Cosmology with Rolling Tachyon

In this section, we study the role of rolling tachyons in the cosmological model with dilatonic gravity. In the string frame, flat space solutions of both initial-stage and late-time will be obtained in closed form. In the Einstein frame, we will show that every expanding solution in flat space is decelerating.

5.1 String frame

We begin with a cosmological model induced from string theory, which is confined on a D3-brane and includes graviton $g_{\mu\nu}$, dilaton Φ , and tachyon T . In the string frame, the effective action of the bosonic sector of the D3-brane system is given by

$$S = \frac{1}{2\kappa^2} \int d^4x \sqrt{-g} e^{-2\Phi} (R + 4\nabla_\mu \Phi \nabla^\mu \Phi) - T_3 \int d^4x e^{-\Phi} V(T) \sqrt{-\det(g_{\mu\nu} + \partial_\mu T \partial_\nu T)}, \quad (74)$$

where we turned off an Abelian gauge field A_μ on the D3-brane and antisymmetric tensor fields of second rank $B_{\mu\nu}$ both on the brane and in the bulk. Equations of motion for the metric, the dilaton, and the tachyon derived from the action (74) are

$$R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} R = \kappa^2 (T_{\mu\nu}^\Phi + e^\Phi T_{\mu\nu}^T), \quad (75)$$

$$4(\nabla_\mu \nabla^\mu \Phi - \nabla_\mu \Phi \nabla^\mu \Phi) + R = \kappa^2 T_3 e^\Phi V(T) \sqrt{1 + \nabla_\alpha T \nabla^\alpha T}, \quad (76)$$

$$\nabla_\mu \nabla^\mu T - \frac{\nabla_\mu \nabla_\nu T \nabla^\mu T \nabla^\nu T}{1 + \nabla_\alpha T \nabla^\alpha T} - \nabla_\mu T \nabla^\mu \Phi - \frac{1}{V} \frac{dV}{dT} = 0, \quad (77)$$

where energy-momentum tensor of the dilaton is

$$T_{\mu\nu}^\Phi = -\frac{2}{\kappa^2} [\nabla_\mu \nabla_\nu \Phi + g_{\mu\nu} (\nabla_\alpha \Phi \nabla^\alpha \Phi - \nabla_\alpha \nabla^\alpha \Phi)], \quad (78)$$

and that of the tachyon

$$T_{\mu\nu}^T = T_3 \frac{V(T)}{\sqrt{1 + \nabla_\alpha T \nabla^\alpha T}} [\nabla_\mu T \nabla_\nu T - g_{\mu\nu} (1 + \nabla_\alpha T \nabla^\alpha T)]. \quad (79)$$

For cosmological solutions in the string frame, we try in addition to the metric ansatz (70)

$$\Phi = \Phi(t). \quad (80)$$

From the action (74), we obtain the following equations

$$3 \left(\frac{\dot{a}^2}{a^2} + \frac{k}{a^2} \right) - 2 \left(3 \frac{\dot{a}}{a} \dot{\Phi} - \dot{\Phi}^2 \right) = \kappa^2 e^\Phi \rho_T, \quad (81)$$

$$2 \frac{\ddot{a}}{a} + \frac{\dot{a}^2}{a^2} + \frac{k}{a^2} - 2 \left(\ddot{\Phi} + 2 \frac{\dot{a}}{a} \dot{\Phi} - \dot{\Phi}^2 \right) = -\kappa^2 e^\Phi p_T, \quad (82)$$

$$4 \left(\ddot{\Phi} + 3 \frac{\dot{a}}{a} \dot{\Phi} - \dot{\Phi}^2 \right) - 6 \left(\frac{\ddot{a}}{a} + \frac{\dot{a}^2}{a^2} + \frac{k}{a^2} \right) = \kappa^2 e^\Phi p_T, \quad (83)$$

$$\frac{\ddot{T}}{1 - \dot{T}^2} + \left(3 \frac{\dot{a}}{a} - \dot{\Phi} \right) \dot{T} + \frac{1}{V} \frac{dV}{dT} = 0, \quad (84)$$

where tachyon energy density ρ_T and pressure p_T defined by $T^{T^\mu}_\nu \equiv \text{diag}(-\rho_T, p_T, p_T, p_T)$ are

$$\rho_T = T_3 \frac{V(T)}{\sqrt{1 - \dot{T}^2}} \quad \text{and} \quad p_T = -T_3 V(T) \sqrt{1 - \dot{T}^2}, \quad (85)$$

which formally coincide with Eq. (20) and Eq. (22). The tachyon equation (84) is equivalent to the following conservation equation

$$\dot{\rho}_T + (3H - \dot{\Phi}) \dot{T}^2 \rho_T = 0, \quad (86)$$

where $H = \dot{a}/a$ is the Hubble parameter. In the absence of detailed knowledge of $V(T)$, we will examine characters of the solutions based on the simplicity of tachyon equation of state

$$p_T = w_T \rho_T, \quad w_T = \dot{T}^2 - 1, \quad (87)$$

which is exactly the same as Eq. (23).

By defining the shifted dilaton $\phi = 2\Phi - 3\ln a$, we rewrite the equations (81)–(84) as

$$\dot{\phi}^2 - 3H^2 + 6\frac{k}{a^2} = 2\kappa^2 e^{\frac{\phi}{2}} a^{\frac{3}{2}} \rho_T, \quad (88)$$

$$2(\dot{H} - H\dot{\phi}) + 4\frac{k}{a^2} = \kappa^2 e^{\frac{\phi}{2}} a^{\frac{3}{2}} p_T, \quad (89)$$

$$\dot{\phi}^2 - 2\ddot{\phi} + 3H^2 + 6\frac{k}{a^2} = -\kappa^2 e^{\frac{\phi}{2}} a^{\frac{3}{2}} p_T, \quad (90)$$

$$\frac{\ddot{T}}{1 - \dot{T}^2} + \frac{1}{2}(3H - \dot{\phi})\dot{T} = -\frac{1}{V} \frac{dV}{dT}. \quad (91)$$

Note that $\sqrt{-g}$, or a^3 is not a scalar quantity even in flat spatial geometry, the shifted dilaton ϕ is not a scalar field in 3+1 dimensions. The conservation equation (86) becomes

$$\dot{\rho}_T + \frac{1}{2}(3H - \dot{\phi})\dot{T}^2 \rho_T = 0. \quad (92)$$

Now Eqs. (88)–(90) and Eq. (87) are summarized by the following two equations

$$2\ddot{\phi} - \dot{\phi}^2 + 2H\dot{\phi} - 3H^2 - 2\dot{H} = -10\frac{k}{a^2}, \quad (93)$$

$$w_T \dot{\phi}^2 + 4H\dot{\phi} - 3w_T H^2 - 4\dot{H} = -(8 + 6w_T)\frac{k}{a^2}. \quad (94)$$

Let us consider only the flat metric ($k = 0$) in the rest part of the paper. If we express the dilaton ϕ as a function of the scale factor $a(t)$, $\phi(t) = \phi(a(t))$, we can introduce a new variable ψ such as

$$\psi \equiv a\phi' = \frac{\dot{\phi}}{H}, \quad (95)$$

where the prime denotes the differentiation with respect to a , and the second equality shows that ψ is the ratio between $\dot{\phi}$ and H . Then Eqs. (93) and (94) are combined into a single first-order differential equation for ψ :

$$4a\psi' + (\psi^2 - 3)(w_T\psi + 2 - w_T) = 0. \quad (96)$$

From now on we look for the solutions of Eq. (96). Above all one may easily find a constant solution $\psi = \mp\sqrt{3}$ which is consistent with Eqs. (88)–(90) only when $\rho_T = 0$:

$$a(t) = a_0(1 \mp \sqrt{3}H_0 t)^{\mp 1/\sqrt{3}}, \quad H(t) = \frac{H_0}{1 \mp \sqrt{3}H_0 t}, \quad (97)$$

$$\Phi(t) = \Phi_0 - \frac{1 \pm \sqrt{3}}{2} \ln(1 \mp \sqrt{3}H_0 t), \quad (98)$$

where $H_0 = H(t=0)$, $a_0 = a(t=0)$, and $\Phi_0 = \Phi(t=0)$ throughout this section. However, exactly-vanishing tachyon density $\rho_T = 0$ from Eq. (88) restricts strictly the validity range of this particular solution to that of vanishing tachyon potential, $V(T) = 0$, which leads to $T = \infty$. The tachyon equation (91) forces $\ddot{T} = 0$ and $\dot{T} = 1$ so that the tachyon decouples ($w_T = p_T = 0$). Therefore, the obtained solution (97)–(98) corresponds to that of string cosmology of the graviton and the dilaton before stabilization but without the tachyon.

Since it is difficult to solve Eq. (96) with dynamical w_T , let us assume that w_T is time-independent (or equivalently $a(t)$ -independent). We can think of the cases where the constant w_T can be a good approximation. From the tachyon potential (8), the first case is onset of tachyon rolling around the maximum point and the second case is late-time rolling at large T region. In fact we can demonstrate that those two cases are the only possibility as far as no singularity evolves.

When w_T is a nonzero constant, Eq. (96) allows a particular solution

$$\psi = \frac{w_T - 2}{w_T} \equiv \beta. \quad (99)$$

This provides a consistent solution of Eqs. (88)–(90)

$$a(t) = a_0 \left(1 + \frac{w_T^2 + 2}{2w_T} H_0 t \right)^{\frac{2w_T}{w_T^2 + 2}}, \quad H(t) = H_0 \left(1 + \frac{w_T^2 + 2}{2w_T} H_0 t \right)^{-1} \quad (100)$$

$$\Phi(t) = \Phi_0 + \frac{2(2w_T - 1)}{w_T^2 + 2} \ln \left(1 + \frac{w_T^2 + 2}{2w_T} H_0 t \right). \quad (101)$$

From Eq. (88) and Eq. (92), the tachyon energy density ρ_T is given by

$$\rho_T(t) = \frac{2 - 2w_T - w_T^2}{w_T^2 \kappa^2 e^{\Phi_0}} H_0^2 \left(1 + \frac{w_T^2 + 2}{2w_T} H_0 t \right)^{-\frac{2(1+w_T)^2}{w_T^2 + 2}}. \quad (102)$$

Since the obtained solution is a constant solution of ψ , it has only three initially-undetermined constants. Specifically, the solution should satisfy $\dot{\Phi} = [(2w_T - 1)/w_T]H$ so that the initial conditions also satisfy a relation $\dot{\Phi}_0 = [(2w_T - 1)/w_T]H_0$. Once we assume general solutions of $a(t)$ -dependent ψ with keeping constant nonzero w_T , they should be classified by four independent parameters ($a_0, H_0, \Phi_0, \dot{\Phi}_0$) instead of three in Eq. (101).

According to the aforementioned condition for valid w_T values, the obtained solution in Eq. (101) may be physically relevant as the onset solution of $w_T = -1$ ($\psi = 3$). In this case, $\rho_T(t)$ is reduced to a constant $\rho_T(t) = 3e^{-\Phi_0} H_0^2 / \kappa^2$. Comparing this with the definition of ρ_T in Eq. (85), the initial Hubble parameter H_0 is related to the dilaton as $H_0 = \pm \kappa e^{\Phi_0/2} \sqrt{T_3/3}$. Then, with $T(t) = 0$, the tachyon equation of motion

is automatically satisfied and hence Eq. (101) becomes an exact solution of the whole set of equations of motion (81)–(84). Since the tachyon field remains as constant at the maximum of the potential, this solution describes the expanding or shrinking solution depending on the initial Hubble parameter, with a constant vacuum energy corresponding to brane tension due to tachyon sitting at the unstable equilibrium point.^b

In order to study the behavior of the tachyon rolling down from the top of the potential, now we slightly perturb this solution, i.e., look for a solution with nonzero but small T dependence. So we treat T as a small expansion parameter and work up to the first-order in T . Since the unperturbed solution satisfies $3H = \dot{\Phi}$, the tachyon equation of motion (84) becomes, to the first-order in T ,

$$\ddot{T} = -\frac{1}{V} \frac{dV}{dT}. \quad (103)$$

This can easily be integrated to

$$\frac{1}{2} \dot{T}^2 = -\ln V + \text{const} = \frac{T^2}{8 \ln 2} + \text{const}, \quad (104)$$

where we used the form of the potential near the origin (8). Given the initial condition that $T = T_0$ and $\dot{T} = 0$ at $t = 0$, we can solve this equation (104) and obtain

$$T(t) = T_0 \cosh \alpha t, \quad (105)$$

where $\alpha = 1/2\sqrt{\ln 2}$. Therefore tachyon starts to roll down the potential as a hyperbolic cosine function. Taking derivative, we find $\dot{T} = \alpha T_0 \sinh \alpha t$. The range for which $\dot{T}$ remains small is then $t \lesssim t_r \equiv 2\sqrt{\ln 2} \sinh^{-1}(2\sqrt{\ln 2}/T_0)$, during which the approximation $w_T \simeq -1$ is good. Unless the initial value T_0 is fine-tuned, the tachyon follows the onset solution (101)–(102) for $t \gtrsim t_r$ and enters into rolling mode.

For more general solutions, the first-order differential equation (96) can be integrated to

$$a = C \left[\frac{\psi^2 - 3}{(\psi - \beta)^2} \left(\frac{\psi - \sqrt{3}}{\psi + \sqrt{3}} \right)^{\frac{\beta}{\sqrt{3}}} \right]^{\frac{\beta-1}{\beta^2-3}}, \quad (106)$$

where C is an integration constant. Note that this algebraic equation does not provide a closed form of ψ in terms of the scale factor $a(t)$ except for a few cases, e.g., $w_T = 0, -1/(\sqrt{3} - 1/2), -2/(3\sqrt{3} - 1)$.

^bThe interpretation as expanding or shrinking solution needs to be more careful, since we are working in the string frame. Actually the behaviors are reversed in the Einstein frame as we will see in the next subsection.

Fortunately, for the late-time case of vanishing w_T , we can obtain the solution in closed form

$$\psi = \sqrt{3} \frac{C_+ \left(\frac{a}{a_0}\right)^{\sqrt{3}/2} + C_- \left(\frac{a}{a_0}\right)^{-\sqrt{3}/2}}{C_+ \left(\frac{a}{a_0}\right)^{\sqrt{3}/2} - C_- \left(\frac{a}{a_0}\right)^{-\sqrt{3}/2}}. \quad (107)$$

Then the scale factor a and the dilaton Φ are explicitly expressed as functions of time t by solving the equations (89)–(90):

$$a(t) = a_0 \left(\frac{C_- t + 2}{C_+ t + 2} \right)^{1/\sqrt{3}}, \quad H(t) = \frac{4H_0}{(C_- t + 2)(C_+ t + 2)}, \quad (108)$$

$$\Phi(t) = \Phi_0 + \ln \left[2 \frac{(C_- t + 2)^{(\sqrt{3}-1)/2}}{(C_+ t + 2)^{(\sqrt{3}+1)/2}} \right], \quad (109)$$

where $C_{\pm} = (3 \mp \sqrt{3})H_0 - 2\dot{\Phi}_0$. We also read the tachyon density ρ_T from Eq. (88)

$$\rho_T = C_+ C_- e^{-\Phi_0} \frac{(C_+ t + 2)^{(\sqrt{3}-1)/2}}{(C_- t + 2)^{(\sqrt{3}+1)/2}}. \quad (110)$$

Note that $C_{\pm}$ should have the same sign from the positivity of the tachyon density (110). Let us first consider that both C_+ and C_- are positive. When $C_- > C_+$ or equivalently $H_0 > 0$, the scale factor a is growing but saturates to a finite value such as $a(\infty) = a_0 (C_-/C_+)^{1/\sqrt{3}}$ in the string frame. When $C_- < C_+$, it decreases. When $C_- = C_+$, $H_0 = 0$ so that the scale factor is a constant, $a(t) = a_0$. For all of the cases, the dilaton Φ approaches negative infinity. Note that $w_T = 0$ means late-time, the tachyon density decreases like $\rho_T \sim 1/t$ as $t \rightarrow \infty$. Consistency check by using Eq. (92) or equivalently by Eq. (91) provides us the expected result, $\dot{T} \rightarrow 1$. If both C_+ and C_- are negative, there appears a singularity at finite time irrespective of relative magnitude of C_+ and C_- .

5.2 Einstein frame

In the previous subsection, it was possible to obtain the cosmological solutions analytically in a few simple but physically meaningful limiting cases. To study the physical implications of what we found, however, we need to work in the Einstein frame. In this subsection, we will convert the cosmological solutions obtained in the string frame to those in the Einstein frame and discuss the physical behaviors.

First of all, let us rewrite the action (74) in the Einstein frame of which metric $g_{\mu\nu}^E$ is related to the string metric by a conformal transformation

$$g_{\mu\nu} = e^{2\Phi} g_{\mu\nu}^E. \quad (111)$$

Note that we abbreviate the superscript E for convenience in what follows. Then the action (74) is changed to

$$S = \frac{1}{2\kappa^2} \int d^4x \sqrt{-g} (R - 2\nabla_\mu \Phi \nabla^\mu \Phi) - T_3 \int d^4x \sqrt{-g} e^{3\Phi} V(T) \sqrt{1 + e^{-2\Phi} \nabla_\mu T \nabla^\mu T}. \quad (112)$$

Equations of motion are

$$R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} R = \kappa^2 (T_{\mu\nu}^\Phi + e^{3\Phi} T_{\mu\nu}^T), \quad (113)$$

$$\nabla_\mu \nabla^\mu \Phi = \frac{\kappa^2}{2} T_3 e^{3\Phi} V(T) \frac{3 + 2e^{-2\Phi} \nabla_\alpha T \nabla^\alpha T}{\sqrt{1 + e^{-2\Phi} \nabla_\alpha T \nabla^\alpha T}}, \quad (114)$$

$$\nabla_\mu \nabla^\mu T + \nabla_\mu T \nabla^\mu \Phi - \frac{e^{-2\Phi} (\nabla_\mu \nabla_\nu T - \nabla_\mu \Phi \nabla_\nu T) \nabla^\mu T \nabla^\nu T}{1 + e^{-2\Phi} \nabla_\alpha T \nabla^\alpha T} - \frac{e^{2\Phi}}{V} \frac{dV}{dT} = 0, \quad (115)$$

where the corresponding energy-momentum tensors are

$$T_{\mu\nu}^\Phi = \frac{1}{\kappa^2} (2\nabla_\mu \Phi \nabla_\nu \Phi - g_{\mu\nu} \nabla_\alpha \Phi \nabla^\alpha \Phi), \quad (116)$$

$$T_{\mu\nu}^T = T_3 \frac{V(T)}{\sqrt{1 + e^{-2\Phi} \nabla_\alpha T \nabla^\alpha T}} [e^{-2\Phi} \nabla_\mu T \nabla_\nu T - g_{\mu\nu} (1 + e^{-2\Phi} \nabla_\alpha T \nabla^\alpha T)]. \quad (117)$$

For cosmological solutions, we use again Eq. (70) and Eq. (80) and then the equations of motion (114)–(115) become

$$\frac{\dot{a}^2}{a^2} + \frac{k}{a^2} = \frac{\kappa^2}{3} (\rho_\Phi + e^{3\Phi} \rho_T) = \frac{1}{3} \dot{\Phi}^2 + \frac{\kappa^2}{3} T_3 e^{3\Phi} V(T) \frac{1}{\sqrt{1 - e^{-2\Phi} \dot{T}^2}}, \quad (118)$$

$$\begin{aligned} \frac{\ddot{a}}{a} &= -\frac{\kappa^2}{6} [(\rho_\Phi + 3p_\Phi) + e^{3\Phi} (\rho_T + 3p_T)] \\ &= -\frac{2}{3} \dot{\Phi}^2 + \frac{\kappa^2}{6} T_3 e^{3\Phi} V(T) \frac{2 - 3e^{-2\Phi} \dot{T}^2}{\sqrt{1 - e^{-2\Phi} \dot{T}^2}}, \end{aligned} \quad (119)$$

$$\ddot{\Phi} + 3\frac{\dot{a}}{a}\dot{\Phi} = -\frac{\kappa^2}{2} e^{3\Phi} (\rho_T - 2p_T) = -\frac{\kappa^2}{2} T_3 e^{3\Phi} V(T) \frac{3 - 2e^{-2\Phi} \dot{T}^2}{\sqrt{1 - e^{-2\Phi} \dot{T}^2}}, \quad (120)$$

$$\frac{\ddot{T}}{1 - e^{-2\Phi}\dot{T}^2} + 3H\dot{T} + \dot{\Phi}\dot{T}\frac{1 - 2e^{-2\Phi}\dot{T}^2}{1 - e^{-2\Phi}\dot{T}^2} + e^{2\Phi}\frac{1}{V}\frac{dV}{dT} = 0. \quad (121)$$

Tachyon energy density ρ_T and pressure p_T in the Einstein frame are obtained by the replacement $\dot{T}_s = e^{-\Phi}\dot{T}$ in Eq. (85),

$$\rho_T = T_3 \frac{V(T)}{\sqrt{1 - e^{-2\Phi}\dot{T}^2}} \quad \text{and} \quad p_T = -T_3 V(T) \sqrt{1 - e^{-2\Phi}\dot{T}^2}, \quad (122)$$

and thereby w_T is given as

$$w_T \equiv p_T/\rho_T = e^{-2\Phi}\dot{T}^2 - 1. \quad (123)$$

Alternatively, we can obtain the same equations from Eqs. (81)–(84) by using the metric of the form ^c

$$ds^2 = e^{2\Phi}(-dt^2 + a^2(t)d\Omega_k^2), \quad (124)$$

and hence the time t and the scale factor a are related to those in the string frame as

$$a_s = ae^{\Phi}, \quad dt_s = e^{\Phi}dt. \quad (125)$$

Then the Einstein equations for the flat case ($k = 0$) in the string frame (81)–(84) are converted to

$$H^2 = \frac{1}{3}\dot{\Phi}^2 + \frac{1}{3}\kappa^2 e^{3\Phi}\rho_T, \quad (126)$$

$$\dot{H} = -\dot{\Phi}^2 - \frac{1}{2}\kappa^2 e^{\Phi}\rho_T\dot{T}^2. \quad (127)$$

Demanding constant w_T is nothing but asking a strong proportionality condition between the dilaton and tachyon, $\dot{T} \propto e^{\Phi}$. Note that the pressure p_T as shown in Eq. (85) is always negative irrespective of both specific form of the tachyon potential ($V(T) \geq 0$) and the value of the kinetic term ($e^{-2\Phi}\dot{T}^2 \leq 1$), and the value of w_T interpolates smoothly between -1 and 0 .

First we observe that the right-hand side of Eq. (126) is always positive, which means that the Hubble parameter $H(t)$ is either positive definite or negative definite for all t and it cannot change the sign in the Einstein frame. Obviously it is a natural consequence of the weak energy condition. Let us first consider the case of positive Hubble parameter, $H(t) > 0$. Eq. (127) shows $\dot{H}$ consists of two terms both of which are negative definite for all t . Since $H > 0$ by assumption, the only consistent behavior of H in this case is that $\dot{H}$ vanishes as $t \rightarrow \infty$, which, in turn implies that $\dot{\Phi}$ and $e^{\Phi}\rho_T\dot{T}^2$ go to zero, separately. It also means that H should be a regular function for

^cIn this subsection all the quantities are in the Einstein frame except the variables with subscript s which denote the quantities in the string frame.

all t . In order to find the large t behavior of $H(t)$, one has to study $e^{-\Phi}\dot{T}$ in large t limit which appears in the definition of w_T in the Einstein frame. Knowing that the functions are regular, it is not difficult to show that the only possible behavior is $e^{-\Phi}\dot{T} \rightarrow 1$ as $t \rightarrow \infty$ after some straightforward analysis of Eqs. (126)–(127). Combining it with the fact that $\dot{\Phi}$ and $e^{\Phi}\rho_T\dot{T}^2$ vanish, we can immediately conclude from Eq. (126) that $H(t)$ should go to zero in large t limit.

The asymptotic behavior of fields in case of the positive Hubble parameter can be found from the solution (109) since w_T is essentially zero for large t as we just have seen above. The only thing to do is to transform the expressions in the string frame to those in the Einstein frame, using the relation (125). Therefore, for large t , we find

$$a(t_s) = a_s(t_s)e^{-\Phi(t_s)} \simeq \frac{1}{2}a_{s0}e^{-\Phi_0}(C_+t_s + 2)^{\frac{\sqrt{3}+1}{2\sqrt{3}}}(C_-t_s + 2)^{\frac{\sqrt{3}-1}{2\sqrt{3}}},$$

$$t = \int dt_s e^{-\Phi} \simeq 2e^{-\Phi_0} \int dt_s \frac{(C_+t_s + 2)^{(\sqrt{3}+1)/2}}{(C_-t_s + 2)^{(\sqrt{3}-1)/2}}. \quad (128)$$

One can also identify the initial Hubble parameter H_0 in terms of $C_{\pm}$ as

$$H_0 = \frac{1}{4}e^{\Phi_0} \left[\left(1 - \frac{1}{\sqrt{3}}\right) C_- + \left(1 + \frac{1}{\sqrt{3}}\right) C_+ \right]. \quad (129)$$

Note that $C_{\pm}$ have the same sign as the Hubble parameter H . Now, with $C_{\pm} > 0$, one can easily confirm from the above equation (128) that all the functions indeed behave regularly. In $t_s \rightarrow \infty$ limit, $a \sim t_s$ and $t \sim t_s^2$ so that the asymptotic behavior of the scale factor becomes $a \sim t^{1/2}$. This power law expansion in flat space is contrasted with the result of Einstein gravity without the dilaton Φ , where ultimately the scale factor ceases to increase, $\lim_{t \rightarrow \infty} a(t) \rightarrow \text{constant}$. The behavior of tachyon density ρ_T can be read from Eq. (110) with t replaced by t_s , which shows that $\rho_T \sim t^{-1/2}$. Since w_T also goes to zero, the fluid of condensed tachyon becomes pressureless. Differently from ordinary scalar matter where matter domination of pressureless gas is achieved for the minimum kinetic energy ($\dot{T} \rightarrow 0$), it is attained for the maximum value of time dependence ($e^{-\Phi}\dot{T} \rightarrow 1$ as $T \rightarrow \infty$) for the tachyon potential given in Eq. (8).

When the Hubble parameter H is negative, the situation is a bit more complicated. Since $\dot{H} < 0$ always, H becomes more and more negative and there is a possibility that eventually H diverges to negative infinity at some finite time. Indeed, it turns out that all solutions in this case develop a singularity at some finite time at which $H \rightarrow -\infty$ and $a \rightarrow 0$. These big crunch solutions may not describe viable universes in the sense of observed cosmological data. Depending on initial conditions, the dilaton Φ diverges to either ∞ or $-\infty$ and $\dot{T}$ goes to either ∞ or zero with the factor $e^{-\Phi}\dot{T}$

remaining finite. It is rather tedious and not much illuminating to show this explicitly, so here we will just content ourselves to present a simple argument to understand the behavior. Since the tachyon field T rolls down from the maximum of the potential to the minimum at infinite T , it is physically clear that $\dot{T}_s = e^{-\Phi}\dot{T}$ would eventually go to one unless there is a singularity at some finite time. Suppose that there appeared no singularity until some long time had passed so that $e^{-\Phi}\dot{T}$ approached to one sufficiently closely. Then Eq. (128) should be a good approximate solution in this case. However, we know that both $C_{\pm}$ are negative when $H < 0$ and Eq. (128) is clearly singular in this case. We have also verified the singular behavior for various initial conditions using numerical analysis.

As mentioned in the previous section, the tachyon T is decoupled when $e^{-\Phi}\dot{T} = 1$ and $T = \infty$. In this decoupling limit, characters of the Einstein equations (126)–(127) that $H^2 > 0$ and $\dot{H} < 0$ do not change so that all the previous arguments can be applied. Well-known cosmological solution of the dilaton gravity before stabilization of the dilaton is

$$a(t) = a_0(1 + 3H_0t)^{1/3}, \quad H(t) = \frac{H_0}{1 + 3H_0t}, \quad (130)$$

$$\Phi(t) = \Phi_0 \pm \frac{1}{\sqrt{3}} \ln(1 + 3H_0t), \quad (131)$$

where the $(\pm)$ sign in Eq. (131) is due to the reflection symmetry ($\Phi \leftrightarrow -\Phi$) in the equations (126)–(127). This solution can also be obtained throughout a transformation (125) from Eqs. (97)–(98). For $H_0 < 0$, it is a big crunch solution ($a \rightarrow 0$) which encounters singularity ($H \rightarrow \infty$, $\Phi \rightarrow \mp\infty$) as $t \rightarrow 1/3|H_0|$. For $H_0 > 0$, it is an expanding but decelerating solution. Since $a \sim t^{1/3}$, the power of expansion rate is increased from $1/3$ to $1/2$ by the tachyonic effect as expected.

So far we discussed generic properties and asymptotic behaviors of solutions in the Einstein frame. Now we consider the behavior at the onset. The solution (101) obtained by assuming constant w_T is transformed to the Einstein frame as

$$\begin{aligned} a(t) &= a_0 \left[1 + \frac{(w_T - 2)^2}{2(1 - w_T)} H_0 t \right]^{\frac{2(1-w_T)}{(w_T-2)^2}}, \\ e^{\Phi(t)} &= e^{\Phi_0} \left[1 + \frac{(w_T - 2)^2}{2(1 - w_T)} H_0 t \right]^{\frac{2(2w_T-1)}{(w_T-2)^2}}, \end{aligned} \quad (132)$$

where the initial Hubble parameter H_0 is related to that in the string frame by $H_0 = e^{\Phi_0} H_{0s} (1 - w_T)/w_T$. Note that H_0 and H_{0s} have opposite signs since $w_T < 0$. Therefore the expanding (shrinking) solution in the string frame corresponds to the shrinking (expanding) solution in the Einstein frame. For

the onset solution with $w_T = -1$, the tachyon energy density ρ_T is a constant as before, $\rho_T(t) = 3e^{-3\Phi_0}H_0^2/4\kappa^2$. Then the initial Hubble parameter is given by $H_0 = \pm 2\kappa e^{3\Phi_0/2}\sqrt{T_3/3}$, which describes the exact solution that tachyon remains at the origin as explained in section 2. Under a small perturbation tachyon starts rolling down according to Eq. (105) with t replaced by t_s . The rest of the discussion on the rolling behavior is the same as in the string frame and the details will not be repeated here.

In conclusion the cosmological solution can be classified into two categories depending on the value of the Hubble parameter $H(t)$ in the Einstein frame. When the initial Hubble parameter H_0 is positive, the solution is regular and the universe is expanding but decelerating as $a(t) \sim \sqrt{t}$ while $e^{\Phi(t)}$ vanishes. When H_0 is negative, there appears a singularity at some finite time t at which the universe shrinks to zero.

6 Summary

In this paper we have discussed basic field theoretic properties of rolling tachyon and Abelian gauge field, and then cosmological solutions. We have shown that both tachyon and Abelian gauge field cannot propagate in terms of perturbative modes when the homogeneous tachyon rolled to the infinity which corresponds to the minimum of its potential. Throughout the study of homogeneous and isotropic universes in string frame, we found initial-stage and late-time solutions in closed form in addition to the known solution of the gravitons and dilaton in the decoupling limit of the tachyon. We have also provided a description of cosmological solutions in the Einstein frame : Once the universe starts expanding, then it continues expanding eternally but decelerating.

We conclude with the list of intriguing questions for further study. The nonlocal effective action we dealt with is not valid for large value of the tachyon field, so genuine string theory description is needed in such regime, including tower of massive string spectra. Since both the tachyon and Abelian gauge field cannot support propagating physical degrees in a perturbative way, an intriguing question may be existence of extended objects in the background of the rolling tachyon configuration. In relation with tachyon cosmology, we may raise a number of questions. Stabilization mechanism of dilaton should be understood and its positive effect to our unsatisfactory decelerating universe solutions is carefully investigated. Various topics for the rolling tachyons asked in the Einstein gravity should be addressed again in the context of dilaton gravity such as existence of inflationary era, possibility as a source of quintessence, reheating without oscillating tachyon modes, cosmological perturbation and structure formation, and so on. When the Abelian gauge field is concerned, one can ask whether or not radiation dominated era is

possible.

Acknowledgments

Three of the authors(C.Kim, H.B. Kim, and Y. Kim) never forget Pong Youl Pac as our teacher and mentor. Y. Kim would like to acknowledge the hospitality of E-Ken researchers of Nagoya University where a part of this work has been done. This work was supported by Korea Research Foundation Grant KRF-2002-070-C00025(C.K.) and the Swiss Science Foundation, grant 21-58947.99(H.B.K.), and is the result of research activities (Astrophysical Research Center for the Structure and Evolution of the Cosmos (ARCSEC) and the Basic Research Program, R01-2000-000-00021-0) supported by Korea Science & Engineering Foundation(Y.K. and O.K.).

References

1. M.R. Garousi, Nucl. Phys. **B584**, 284 (2000), hep-th/0003122; E.A. Bergshoeff, M. de Roo, T.C. de Wit, E. Eyras, and S. Panda, JHEP **0005**, 009 (2000), hep-th/0003221; J. Klusen, Phys. Rev. **D62**, 126003 (2000), hep-th/0004106.
2. S. Sugimoto and S. Terashima, JHEP **0207**, 025 (2002), hep-th/0205085; J.A. Minahan, JHEP **0207**, 030 (2002), hep-th/0205098.
3. G.W. Gibbons, K. Hori, and P. Yi, Nucl. Phys. **B596**, 136 (2001), hep-th/0009061.
4. A.A. Gerasimov and S.L. Shatashvili, JHEP **0010**, 034 (2000), hep-th/0009103; D. Kutasov, M. Marino, and G.W. Moore, JHEP **0010**, 045 (2000), hep-th/0009148; hep-th/0010108.
5. A. Sen, JHEP **0204**, 048 (2002), hep-th/0203211; hep-th/0203265; Mod. Phys. Lett. **A17**, 1797 (2002), hep-th/0204143; hep-th/0207105; hep-th/0209122; N. Moeller and B. Zwiebach, hep-th/0207107.
6. A. Ishida and S. Uehara, Phys. Lett. **B544**, 353 (2002), P. Mukhopadhyay and A. Sen, hep-th/0208142; hep-th/0206102; G. Gibbons, K. Hashimoto, and P. Yi, hep-th/0209034.
7. T. Mehen and B. Wecht, hep-th/0206212.
8. S.-J. Rey and S. Sugimoto, hep-th/0301049.
9. C. Kim, H.B. Kim, Y. Kim, and O-K. Kwon, hep-th/0301076.
10. For a review of the string cosmology, see one of the followings : G. Veneziano, 628p in the proceedings of Les Houches 1999, (Berlin, Springer-Verlag, 2001), hep-th/0002094; N.E. Mavromatos, hep-th/0111275; E.J. Copeland, hep-th/0202028.
11. G. W. Gibbons, Phys. Lett. **B537**, 1 (2002), hep-th/0204008; For a review, see hep-th/0301117.

12. S. Alexander, Phys. Rev. D**65**, 023507 (2002), hep-th/0105032; A. Mazumdar, S. Panda, and A. Perez-Lorezana, Nucl. Phys. B**614**, 101 (2001), hep-ph/0107058.
13. L. Kofman and A. Linde, JHEP **0207**, 004 (2002), hep-th/0205121; A.V. Frolov, L. Kofman, and A.A. Starobinsky, Phys. Lett. B**545**, 8 (2002), hep-th/0204187; G.N. Felder, L. Kofman, and A.A. Starobinsky, JHEP **0209**, 026 (2002), hep-th/0208019.
14. M. Fairbairn and M.H.G. Tytgat, hep-th/0204070; S. Mukohyama, Phys. Rev. D**66**, 024009 (2002), hep-th/0204084; A. Feinstein, Phys. Rev. D**66**, 063511(2002), hep-th/0204140; T. Padmanabhan, Phys. Rev. D**66**, 021301(R) (2002), hep-th/0204150; D. Choudhury, D. Ghoshal, D.P. Jatkar, and S. Panda, Phys. Lett. B**544**, 231 (2002), hep-th/0204204; X. Li, J. Hao and D. Liu, Chin. Phys. Lett. **19**, 1584 (2002), hep-th/0204252; G. Shiu and I. Wasserman, Phys. Lett. B**541**, 6 (2002), hep-th/0205003; H.B. Benaoum, hep-th/0205140; M. Sami, hep-th/0205146; M. Sami, P. Chingangbam, and T. Qureshi, Phys. Rev. D**66**, 043530 (2002), hep-th/0205179; J. Hwang and H. Noh, hep-th/0206100; Y.-S. Piao, R.-G. Cai, X. Zhang, and Y.-Z. Zhang, hep-ph/0207143; G. Shiu, S.-H. H. Tye, and I. Wasserman, hep-th/0207119; X.-z. Li, D.-j. Liu, and J.-g. Hao, hep-th/0207146; J.M. Cline, H. Firouzjahi, and P. Martineau, hep-th/0207156; S. Mukohyama, hep-th/0208094; M.C. Bento, O. Bertolami, and A.A. Sen, hep-th/0208124; J.-g. Hao and X.-z. Li, Phys. Rev. D, hep-th/0209041; J.S. Bagla, H.K. Jassal, and T. Padmanabhan, astro-ph/0212198.
15. C. Kim, H.B. Kim, and Y. Kim, Phys. Lett. B**552**, 111 (2003), hep-th/0210101.
16. For example, see Chapter 3 and 8 of *String Theory* by J. Polchinski, (Cambridge University Press, 1998).